


Book of Abstracts



**9th Conference of the
European Foundation for Plant Pathology**



**6th Congress of the
Sociedade Portuguesa de Fitopatologia**



**Integrated Plant Disease
Management**



Évora, Portugal

Nov 15-18, 2010

**9TH CONFERENCE OF THE EUROPEAN FOUNDATION FOR PLANT PATHOLOGY
AND 6TH CONGRESS OF THE SOCIEDADE PORTUGUESA DE FITOPATOLOGIA**

President

Maria Ivone Clara (Universidade de Évora)

Scientific Committee

Alan Phillips (Universidade Nova de Lisboa)

Maria Ivone Clara (Universidade de Évora)

Gustavo Nolasco (Universidade do Algarve)

Manuel Mota (Universidade de Évora)

Pedro Talhinhos (Instituto de Investigação Científica e Tropical)

Marta Laranjo (Universidade de Évora)

Local Organizing Committee (Universidade de Évora)

Maria Ivone Clara

Maria do Rosário Félix

Carla Marisa Varanda

Marta Laranjo

José Calado

Fernando Rei

Ana Alexandre

Manuel Guerreiro

Maria Mário Azedo



INTRODUCTION

In recent years, a novel class of necrosis-inducing proteins, known as Nep1-like proteins (NLPs) has been identified in bacteria, fungi and oomycetes. Species of the genus *Phytophthora* are causal agents of serious plant diseases infecting *Castanea sativa*. *Phytophthora cinnamomi* secrete necrosis-inducing protein (NPP1) causing necrosis on leaf and roots of the plant, leading to death. The *NPP1* ORF contains 771 bp and encodes a 256 aa protein. The study of factors that affect *NPP1* gene expression is extremely important to better evaluate the mechanism of plant necrosis induced by *Phytophthora cinnamomi*. In order to understand its function, we evaluated the expression in different inducers media by RT-qPCR. The medium rich in glucose as carbon source were the one who showed major levels of expression.

MATERIALS AND METHODS

Genomic DNA was obtained from strain *P. cinnamomi* Pr120 as described by Cooke and Duncan (1997). Polymerase chain reaction was used to amplify a fragment of the *NPP1* gene, using degenerate oligonucleotide which were designed based on homology of previous published *Phytophthora* sp. NPP's sequences from EMBL databases. PCR was performed with 10X amplification buffer; 100ng DNA template; 0.2 mM dATP, dCTP, dGTP, and dTTP (each); 0.2µM each primer; 1.5mM MgCl2; and 1U Taq DNA polymerase (Promega), in a final reaction volume of 50 µl. Amplification consisted of: one cycle of 5 min at 94°C, and 36 cycles of denaturing for 1 min at 94°C; annealing for 1 min at 63°C; and extension for 30 s at 72°C. A final extension step of 5 min at 72°C was done for one cycle. Full gene sequence length elucidation (1328bp) was achieved by flanking the previous known sequence with asymmetric HE-TAIL PCR using the degenerated primers R1, R2, R3 and R4 and the methodology described by Michiels *et al.*(2003), and the gene-specific primers. Three rounds of PCR were performed on a MyCycler Thermal Cycler (BIORAD), using the product of the previous PCR as a template for the next. A detailed cycler program and conditions are given in Table1.

DNA sequencing was performed using an ABI 373 automated sequencer. The open reading frame (ORF) of *P. cinnamomi* *NPP1* was identified by BioEdit program and submitted to EMBL databases (Accession number AM403130); Nucleotide and amino acid sequences were analyzed using FASTA programs from EMBL databases. ClustalW2 (Larkin *et al.*, 2007) was used to align the *Phytophthora* genus NPP1 sequences..

Gene Expression: The expression was studied during growth in different carbon sources and was also performed a time course of *NPP1* production. To analyze gene expression was performed RT-qPCR using the *SyBr Green* method and the actin gene *Act2* as a endogenous control.

RESULTS AND DISCUSSION

The translated ORF of *P. cinnamomi* *NPP1* codifies a 256 aa protein, with a predict Mw of 29 KDa and a theoretical iso-electric point of 8,58.

In Figure 2 are shown the multiple alignment of various sequences who showed great homology with *P. cinnamomi* *NPP1*, including another *NPP*'s of *Phytophthora* genus, and a *NPP1* super-family conserved domain from oomycetes.




FIGURE 2 – Multiple sequence alignment of *NPP1* genes from *Phytophthora* sp.

TABLE 1 - HE-TAIL PCR cycle settings and conditions*

Reaction	Number of Cycles	Thermal Settings
Prim ary	1	93°C (1min); 95°C (5min)
	5	94°C (30seg), 62°C (1min), 72°C (2min30s)
	1	94°C (30seg), 25°C ramping 72°C (3min); 72°C (2min30s)
	15	94°C (20s), 65°C (3min30s); 94°C (20s), 65°C (3min30s); 94°C (30s), 42°C (1min), 72°C (2min30s) 72°C (5min), 4°C Hold
Secondary	1	94°C (20s); 65°C (3min30s); 94°C (20s); 65°C (3min30s); 94°C (30s); 42°C (1min); 72°C (2min30s) 72°C (5min), 4°C Hold
	12	94°C (20s); 65°C (3min30s); 94°C (20s); 65°C (3min30s); 94°C (30s); 42°C (1min); 72°C (2min30s) 72°C (5min), 4°C Hold
	1	94°C (30s); 42°C (1min); 72°C (2min30s) 72°C (5min), 4°C Hold
Tertiary	30	94°C (30s); 42°C (1min); 72°C (2min30s)
	1	72°C (5min); 4°C Hold




FIGURE 1 - RT-qPCR of *NPP1* gene from *P. cinnamomi*.

The study of factors that affect *NPP1* gene expression is extremely important to better evaluate the mechanism of plant necrosis induced by *Phytophthora cinnamomi*. In order to understand its function, we evaluated the expression in different inducers media by RT-qPCR . The medium rich in glucose as carbon source were the one who showed major levels of expression in Figure 1.

REFERENCES

Cooke, D. & Duncan, J., 1997. Phylogenetic analysis of *Phytophthora* species based on the ITS1 and ITS2 sequences of ribosomal DNA. *Mycological Research*, 101:667-677.
Larkin, M., Blackshields, G., Brown, N., Chenna, R., McGettigan, P., McWilliam, H., Valentin, F., Wallace, I., Wilm, A., Lopez, R., Thompson, J., Gibson, T. & Higgins D., 2007. Bioinformatics, 23(21): 2947-2948.
Michiels, A., Tucker, M., Van Den Ende, W. & Van Laere, A., 2003. Chromosomal Walking of Flanking Regions From Short Known Sequences in GC-Rich Plant Genomic DNA. *Plant Molecular Biology Reporter*, 21:295-302.
Rose, J., Ham, K., Darvill, A. & Albersheim, P., 2002. Molecular cloning and characterization of glucanase inhibitor proteins: coevolution of a counterdefense mechanism by plant pathogens. *Plant Cell*, 14:1329-1345.

Acknowledgements:

The Project COMBATINTA/SP2.P11/02 Interreg IIIA – Cross-Border Cooperation Spain-Portugal, financed by The European Regional Development Fund, and the Project “Identification, characterization and role of molecular factors associated with the mechanisms of infection of Fagaceae species by *Phytophthora cinnamomi*” (PTDC/AGR-AAM/67628/2006) FCT, supported this work.